

CONTENTS

| | |
|--|-------|
| Potential health benefits of major seed spices – S. S. Rathore, S. N. Saxena and Balraj Singh | 1-12 |
| Effect of moisture content, screw and grinder speed on physical and thermal properties of fenugreek powder – P. Barnwal, K. K. Singh, Alka Sharma, A. K. Choudhary, Chothani Hardik R, S.N.Saxena | 13-21 |
| Genetic Variability and Character Association in Fenugreek (<i>Trigonella foenum- graecum</i> L.) – Anubha Jain, Balraj Singh, R. K. Solanki, S.N. Saxena and R.K. Kakani | 22-28 |
| Effect of irrigation and different seed treatment methods on growth and yield of fenugreek (<i>Trigonella foenum graecum</i> L.) – G. Lal, I. P. Saini, R.S. Mehta, S.P. Maheria and Yash Sharma | 29-33 |
| Effect of foliar application of plant growth regulators on growth, yield and essential oil components of Ajwain (<i>Trachyspermum ammi</i> L.) – Yogita Rohamare, T. D. Nikam and K. N. Dhumal | 34-41 |
| Effect of bioregulators on productivity and profitability of fenugreek (<i>Trigonella foenum-graecum</i> L.) – A.S. Godara, U.S. Gupta, Ravindra Singh and G.Lal | 42-45 |
| Prevalence of cumin diseases on farmer's field: A survey of Rajasthan and Gujarat states – Y. K. Sharma, K. Kant, R. K. Solanki and R. P. Saxena | 46-49 |
| Relative abundances and foraging behaviour of honey bee species on minor seed spice crops – Krishna Kant, Balraj Singh, S. R. Meena, J. K. Ranjan, B. K. Mishra, R.K. Solanki and Mukesh Kumar | 50-53 |
| Effect different sources of organic manures with and without bio fertilizers in Cumin (<i>Cuminum cyminum</i> L.) – S. G. Patel, A. U. Amin, S. P. Patel, A. V. Agalodiya and S. M. Patel | 54-58 |
| Microbiological profile of coriander (<i>Coriandrum sativum</i> L.) crop rhizosphere in Rajasthan and screening for auxin producing rhizobacteria – B. K. Mishra, Ashutosh Sharma, O.P. Aishwath, Y.K. Sharma, Krishna Kant, M.K. Vishal, S.N. Saxena and J.K. Ranjan | 59-64 |
| Effect of technological interventions on cumin yield at farmers' field – G. Lal ¹ , R.S. Mehta ¹ , D. Singh and M.K. Choudhary | 65-69 |
| Effect of phosphorus and sulphur levels on the performance of Nagauri Methi (<i>Trigonella corniculata</i> L.) under semi arid areas of Rajasthan – A.S. Godara, S. Kapade, G. Lal and Ravindra Singh | 70-73 |
| Nutrients uptake pattern in some important cultivars of cumin for nutritional budgeting – O.P. Aishwath and S.K. Malhotra | 74-80 |
| Effect of sowing time and crop geometry on growth and seed yield of dill (<i>Anethum sowa</i> L.) – S.S. Meena, R.S. Mehta, R.D. Meena, N. K. Meena and B. Singh | 81-84 |
| Physiological parameters and their relation to seed yield in coriander (<i>Coriandrum sativum</i> L.) – S. N. Saxena, S. S. Rathore, Rohit Saxena and R. K. Kakani | 85-88 |
| Performance of coriander (<i>Coriandrum sativum</i> L.) varieties for growth and seed yield – T.P. Malik and S. K. Tehlan | 89-90 |
| Rapid and mass screening method for galactomannan content in fenugreek seeds – S.S.Rathore, S.N.Saxena, R. K. Kakani and Balraj Singh | 91-93 |
| Seed spices - a better option for crop diversification in Nagaur district of Rajasthan – O.P. Garhwal, Dinesh Arora, S.K Sharma, Ramankant Sharma and R.Porwal | 94-96 |